

Does the Fine Structure Constant Really Vary in Time?

Mario Livio and Massimo Stiavelli^{1,2}

Space Telescope Science Institute

3700 San Martin Drive

Baltimore, MD 21218

Received _____; accepted _____

¹On leave from the Scuola Normale Superiore, Piazza dei Cavalieri 7, I56126 Pisa, Italy

²On assignment from the Space Science Dept. of the European Space Agency

ABSTRACT

We discuss how laboratory experiments can be used to place constraints on possible variations of the fine structure constant α in the observationally relevant redshift interval $z \simeq 0 - 5$, within a rather general theoretical framework. We find a worst case upper limit for $\Delta\alpha/\alpha$ of 8×10^{-6} for $z \leq 5$ and $\Delta\alpha/\alpha$ of 0.9×10^{-6} for $z \leq 1.6$. The derived limits are at variance with the recent findings by Webb et al. (1998), who claim an observed variation of $\Delta\alpha/\alpha = -2.6 \pm 0.4 \times 10^{-5}$ at $1 < z < 1.6$.

subject headings: cosmology: theory - physical data and theory: relativity, nuclear reactions, nucleosynthesis, abundances

1. Introduction

In an interesting paper, Webb et al. (1998) describe an observational method for investigating possible time and space variations in the fine structure constant α , using quasar absorption systems (see also Drinkwater et al. 1998). Webb et al. present intriguing evidence suggesting that α was smaller in the past (for redshifts $z > 1$). The claimed fractional change is (for $z > 1$), $\Delta\alpha/\alpha = -2.6 \pm 0.4 \times 10^{-5}$.

Since a varying value of α can, in principle at least, have significant consequences for cosmology (e.g. concerning recombination), we examine in the present letter possible variations in α in a broader context, although still under a particular set of assumptions. We also show that in the context of the theory presented in this paper the result obtained by Webb et al. appears to be in conflict with results of tests of the equivalence principle.

2. By How Much Can α Change?

The experimental constraints on variations of α were explored extensively by Bekenstein (1982) by constructing a dynamical theory relating variations of α to the electromagnetic fraction of the mass density in the universe. Bekenstein’s (1982) framework for α variability was based on *very general assumptions*: covariance, gauge invariance, causality, time reversal of electromagnetism, and that gravitation is described by a metric of spacetime which satisfies Einstein’s equations. He obtained the following equation for the temporal variation of α (adopting a Robertson-Walker metric; we correct here a misprint in the original paper)

$$(a^3 \dot{\epsilon}/\epsilon) = -a^3 \zeta (l^2/\hbar c) \rho_m c^4 \quad . \quad (1)$$

Here $\epsilon = (\alpha/\alpha_{\text{today}})^{1/2}$, l is a length scale of the theory, ρ_m is the total rest mass density of matter, a is the expansion scale factor, and $\zeta = \langle m_{n,em} \rangle / m_p$ is a dimensionless parameter

which measures the fraction of mass in the form of Coulomb energy of an average nucleon, $\langle m_{n,em} \rangle$, compared to the free proton mass, m_p . In order to be able to integrate eq. (1), Bekenstein (1982) assumed that ζ (which is affected by transformations of hydrogen into heavier nuclei in stars) is constant. He was able to show that this assumption is reasonable up to redshifts $z \lesssim 1$, by the following consideration. The proton conversion rate (via the reaction $4\text{H}^1 \rightarrow \text{He}^4$) can be estimated from the mean luminosity density (e.g. Davis, Geller & Huchra 1978) to be $\sim 8 \times 10^{-28} \text{ cm}^{-3} \text{ s}^{-1}$. Since a minimal nucleon number density in the universe can be determined from nucleosynthesis to be (e.g. Olive, Schramm & Steigman 1981) $\sim 10^{-7} \text{ cm}^{-3}$, the timescale for proton conversion (and thereby for a change in ζ) is $\tau_{\text{conv}} \gtrsim 10^{20} \text{ s}$.

The timescale for changes in ρ_m or a is $\sim H_o^{-1} \sim 10^{17} \text{ s}$. Thus, for as far back as the luminosity of galaxies does not change significantly (up to $z \lesssim 1$), the assumption of a constant ζ is a reasonable one.

However, it is interesting to push this analysis to higher redshifts (e.g. the main change in α claimed by Webb et al. (1998) occurs at $z \gtrsim 1$) by extending Bekenstein's (1982) analysis with a determination of the behavior of ζ at higher redshifts. We achieve this as follows. As explained above, the rate of change of the Coulomb contribution to the baryon mass depends on the rate of conversion of hydrogen into helium. Following Bekenstein we find:

$$\zeta \simeq 1.2 \times 10^{-2} (X + \frac{4}{3}Y) \quad , \quad (2)$$

where X and Y are the mass fractions in hydrogen and helium, respectively. In order to determine the conversion rate we adopted three possible star formation rate (SFR) histories for the universe: (i) one that follows the Madau et al. (1996) curve, (ii) one that rises from the present up to a peak at $z \sim 1$ and then is flat at the peak value up to $z \sim 5$ (as is perhaps suggested by the COBE Diffuse Infrared Background experiment; Hauser et al.

1998; Calzetti & Heckman 1998), and (iii) a SFR that is constant (at the peak value) for all redshifts (as a limiting case). These star formation histories were used to synthesize a mean stellar population of the universe as a function of time, using the Bruzual and Charlot (1993) evolutionary models. The bolometric luminosity from the mean stellar population was then used to calculate the hydrogen to helium conversion rate. In Fig. 1, we show the proton conversion rate as a function of redshift up to $z = 5$, for the three assumed SFR histories. As can be seen from the figure, the timescale for proton conversion satisfies $\tau_{\text{conv}} \gtrsim 10^{19} \text{s}$ up to $z = 5$. Since $\tau_{\text{conv}} \gg H_o^{-1}$, we can still take ζ to be constant in integrating eq. (1). Indeed direct calculation of eq. (2) shows that it varies by less than 1 % over the redshift interval that we consider. Thus we obtain:

$$\dot{\epsilon}/\epsilon = -\zeta(l^2 c^3/\hbar)\rho_m(t - t_c) \quad , \quad (3)$$

where t_c is an (unknown) integration constant. From eq. (2) with $X = 0.74$ and $Y = 0.26$ we find for ζ the value 1.3×10^{-2} (see also Bekenstein 1982). We will consider in the following the case of a low Ω ($\Omega \simeq 0.2$, see below) Universe. The results however would not change significantly in a $\Omega = 1$ Universe. Eq. (3) can be integrated in time by taking into account that for $z \ll 4(\Omega^{-1} - 1)$ one has $\rho_m \simeq \Omega_{mo}\rho_c(t_o/t)^3$, where Ω_{mo} is the baryon fraction at the present time t_o , and ρ_c is the critical density $\rho_c \equiv 3H_o^2/(8\pi G)$, and that $|\epsilon - 1| \ll 1$. We find:

$$\epsilon(z) - 1 \simeq -\frac{3\zeta}{8\pi} \left(\frac{l}{L_p} \right)^2 \Omega_{mo} H_o^2 t_o^2 \left(1 + \frac{t_c t_o}{2t^2(z)} - \frac{t_o}{t(z)} - \frac{t_c}{2t_o} \right) \quad , \quad (4)$$

where $L_p = (G\hbar/c^3)^{1/2}$ is the Planck length. We have verified that the above analytical approximation is applicable to the redshift interval of interest. As an example, in Fig. 2 we plot for the case $t_c = t_o$ the difference between the expression of eq. (4) and the result of a direct numerical integration of eq. (3). Eq. (4) has two quite different regimes depending on the value of t_c/t_o . Let us consider first the case $|t_c| \lesssim t_o$ which is perhaps the most

physically founded since there is no a priori reason for having $|t_c| \gg t_o$. We find:

$$|\epsilon(z \leq 5) - 1| \lesssim 6.2 \times 10^{-3} \Omega_{mo} \left(\frac{l}{L_p} \right)^2 . \quad (5)$$

In order to obtain an absolute upper limit on the variability, we will assume that *all* the matter density is nucleonic, by taking $\Omega = \Omega_{mo} \simeq 0.2$ (e.g. Garavich et al. 1998; Perlmutter et al. 1998). We can also use the results of the Eötvös-Dicke-Braginsky experiments (Eötvös, Parker & Fekete 1922; Roll, Krotkov & Dicke 1964; Braginsky & Pamov 1972), designed to test the equivalence principle, to infer $l/L_p \lesssim 10^{-3}$. Thus, our final upper limit is:

$$|\epsilon(z \leq 5) - 1| \lesssim 1.2 \times 10^{-9} . \quad (6)$$

If we assume instead that $|t_c| \gg t_o$ we can set an upper limit to $|t_c|$ by considering constraints on $|\dot{\epsilon}/\epsilon|$. From Shlyakhter's (1976) analysis of the Oklo natural reactor we have $|\dot{\epsilon}/\epsilon| < 0.5 \times 10^{-7} H_o$. More recently, Damour and Dyson (1996) have derived the less strong but more robust limit of $|\dot{\epsilon}/\epsilon| < 3.4 \times 10^{-7} H_o$, which we will adopt in the following. When used in combination with eq. (3) and with $\Omega_{mo} = 0.2$ this gives:

$$|(l/L_p)^2 t_c H_o| < 6.6 \times 10^{-3} . \quad (7)$$

The term involving t_c in Eq. (4) becomes dominant and we obtain:

$$|\epsilon(z \leq 5) - 1| \lesssim 8 \times 10^{-6} . \quad (8)$$

The above limits become stronger as the redshift decreases. For example, at the redshift of interest for the Webb et al. (1998) result ($z=1.6$) one would find instead $|\epsilon(z = 1.6) - 1| \lesssim 1.6 \times 10^{-10}$ if $|t_c|/t_o \lesssim 1$ or $|\epsilon(z = 1.6) - 1| \lesssim 9 \times 10^{-7}$ otherwise. Both limits, regardless of the value of t_c , are at variance with the Webb et al. (1998) result.

3. Discussion

The analysis of experimental constraints for the variation of the fine structure constant within the framework of rather general dynamical theories provides upper limits to any change in α which are at variance with the Webb et. al (1998) detection. We have verified that varying the cosmological parameters (i.e. the values of Ω , H_0 , primordial helium fraction) does not affect our constraints significantly.

We should note that variations in α in the context of the theory discussed in the present paper result in no modifications to the Planckian spectrum of black body radiation. In fact one could think of varying α as a varying permittivity of the vacuum. This would simply be equivalent to changing the expansion factor $a(t)$ to a different function, which cannot modify the Planck spectrum.

Variations of the fine structure constant α could be driven also by a different dynamics. In theories with extra dimensions (e.g., Kaluza-Klein or superstrings theories) variations in the observed value of α can be induced by variations in the length scale of the compact dimensions (see, e.g., Marciano 1984, Barrow 1987). The effects of extra dimensions were discussed in detail by Barrow (1987) who found upper limits to $|\dot{\alpha}/\alpha|$ of $\lesssim 10^{-8}H_0$ for both Kaluza-Klein and superstring theories. This upper limit is also much smaller than the claimed detection by Webb et. al (1998). However, note that in these theories the variations of all physical constants are inter-related.

Finally, we would like to mention that the calculations leading to eq. (4) would have to be modified if a significant component of the dark matter in the universe were to decay electromagnetically at redshift $z \gtrsim 1$ (e.g. Sciama 1982, 1997). We hope that the present letter will inspire more attempts to determine possible variations of α observationally.

ML acknowledges support from NASA Grant NAG5-6857. We thank Yacov Bekenstein

for very useful discussions and an anonymous referee for a careful reading of the manuscript.

REFERENCES

- Barrow J.D., 1987, Phys. Rev. D, 35, 1805
- Bekenstein J.D., 1982 Phys. Rev. D, 25, 1527.
- Braginsky V.B., Panov V.I., 1972, Zh. Eksp. Teor. Fiz. 61, 873 [Sov. Phys.-JETP 34, 463 (1972)]
- Bruzual G., Charlot S., 1993, ApJ 405, 538
- Eötvös R.V., Pekar V., Fekete E., 1922, Ann. Phys. (Leipzig) 68, 11
- Calzetti D., Heckman T.M., 1998, astro-ph/9806083 (ApJ submitted)
- Damour T., Dyson F., 1996, Nucl. Phys., B480, 37
- Davis M., Geller M.J., Huchra J., 1978, ApJ, 221, 1
- Drinkwater M.J., Webb J.K., Barrow J.D., Flambaum V.V., 1998, MNRAS, 295, 457
- Garnavich P.M., et. al , 1998, ApJ, 493L
- Hauser M.G., et al. 1998, ApJ, in press
- Madau P., Ferguson H.C., Dickinson M.E., Giavalisco M., Steidel C.S., Fruchter A., 1996, MNRAS, 283, 1388
- Marciano W., 1984, Phys. Rev. Lett., 52, 1984
- Olive K.A., Schramm D.N., Steigman G., Turner M.S., Yang J., 1981, ApJ. 264, 557.
- Perlmutter S., et. al , 1998, Nature, 391, 51
- Roll P.G., Krotkov R., Dicke R.H., 1964, Ann. Phys. (NY) 26, 442

Sciama D.W. 1982, MNRAS, 198, 1

Sciama D.W. 1997, MNRAS, 289, 945

Shlyakhter A.I., 1976, Nature, 264, 340

Webb J. K., Flambaum V.V., Churchill C.W., Drinkwater M.J., Barrow J.D., 1998,
astro-ph/9803165

Fig. 1.— We plot the proton conversion rate as a function of redshift for a constant SF model (solid line), the Madau et. al cosmic SFR history (dotted line) and a SFR history rising up to $z = 1$ and then staying constant at the peak value (dashed line).

Fig. 2.— Difference between the analytical approximation and the numerically integrated value of ϵ . For redshift $z \leq 10$ the difference between the two is negligible.



